



DOI: 10.4274/jcrpe.galenos.2026.2025-12-10

J Clin Res Pediatr Endocrinol 2026;18(3):459-466

Mucocutaneous Findings and Endocrinopathies in Children with Turner Syndrome: A Cross-Sectional Study

İ Selami Aykut Temiz¹, İ Filiz Cebeci², İ Aslıhan Çiçekli¹, İ Begüm Işık³, İ Sevil Savaş Erdoğan⁴,
İ Selma Emre⁵, İ Hilal Kaya Erdoğan⁶, İ İsa An⁷, İ Sezgi Sarıkaya Solak⁸, İ Sevilay Kılıç⁹,
İ Eda Öksüm Solak¹⁰, İ Arzu Ataseven¹, İ Beray Selver Eklioğlu¹¹, İ Suna Kılınç¹², İ Mehmet Boyraz¹³,
İ Enver Şimşek¹⁴, İ Tuğba Çetin Kontbay¹⁵, İ Filiz Tütüncüler¹⁶, İ Durmuş Doğan¹⁷, İ Ülkü Gül Şiraz¹⁸,
İ Güllü Eren¹⁹

¹Necmettin Erbakan University Faculty of Medicine, Department of Dermatology, Konya, Türkiye

²University of Health Sciences Türkiye, Kartal Dr. Lütfi Kırdar City Hospital, Clinic of Dermatology, İstanbul, Türkiye

³University of Health Sciences Türkiye, Yozgat City Hospital, Clinic of Dermatology, Yozgat, Türkiye

⁴University of Health Sciences Türkiye, Sultan 2. Abdulhamid Han Training and Research Hospital, Clinic of Dermatology, İstanbul, Türkiye

⁵Ankara Yıldırım Beyazıt University Faculty of Medicine, Department of Dermatology, Ankara, Türkiye

⁶Eskişehir Osmangazi University Faculty of Medicine, Department of Dermatology, Eskişehir, Türkiye

⁷Şanlıurfa Training and Research Hospital, Department of Dermatology, Şanlıurfa, Türkiye

⁸Trakya University Faculty of Medicine, Department of Dermatology, Edirne, Türkiye

⁹Çanakkale Onsekiz Mart University Faculty of Medicine, Department of Dermatology, Çanakkale, Türkiye

¹⁰Erciyes University Faculty of Medicine, Department of Dermatology, Kayseri, Türkiye

¹¹Necmettin Erbakan University Faculty of Medicine, Department of Pediatric Endocrinology, Konya, Türkiye

¹²University of Health Sciences Türkiye, Zeynep Kamil Women and Children Diseases Training and Research Hospital, Clinic of Pediatric Endocrinology, İstanbul, Türkiye

¹³Ankara Yıldırım Beyazıt University Faculty of Medicine, Department of Pediatric Endocrinology, Ankara, Türkiye

¹⁴Eskişehir Osmangazi University Faculty of Medicine, Department of Pediatric Endocrinology, Eskişehir, Türkiye

¹⁵Şanlıurfa Training and Research Hospital, Clinic of Pediatric Endocrinology, Şanlıurfa, Türkiye

¹⁶Trakya University Faculty of Medicine, Department of Pediatric Endocrinology, Edirne, Türkiye

¹⁷Çanakkale Onsekiz Mart University Faculty of Medicine, Department of Pediatric Endocrinology, Çanakkale, Türkiye

¹⁸Erciyes University Faculty of Medicine, Department of Pediatric Endocrinology, Kayseri, Türkiye

¹⁹Necmettin Erbakan University Faculty of Medicine, Department of Public Health, Konya, Türkiye

Cite this article as: Temiz SA, Cebeci F, Çiçekli A, Işık B, Savaş Erdoğan S, Emre S, Kaya Erdoğan H, An İ, Sarıkaya Solak S, Kılıç S, Öksüm Solak E, Ataseven A, Eklioğlu BS, Kılınç S, Boyraz M, Şimşek E, Çetin Kontbay T, Tütüncüler F, Doğan D, Şiraz ÜG, Eren G. Mucocutaneous findings and endocrinopathies in children with Turner syndrome: a cross-sectional study. J Clin Res Pediatr Endocrinol. [Epub Ahead of Print] 2026;18(3):459-466

Corresponding Author: Assoc. Prof. Selami Aykut Temiz, Necmettin Erbakan University Faculty of Medicine, Department of Dermatology, Konya, Türkiye

E-mail: aykutmd42@gmail.com **ORCID:** orcid.org/0000-0003-4878-0045

Conflict of interest: None declared

Received: 18.12.2025 **Accepted:** 24.02.2025 **Epub:** 26.02.2026 **Publication Date:** 08.09.2026



©Copyright 2026 by Turkish Society for Pediatric Endocrinology and Diabetes / The Journal of Clinical Research in Pediatric Endocrinology published by Galenos Publishing House.
Licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 (CC BY-NC-ND) International License.

ABSTRACT

Objective: To investigate the frequency of skin, hair, nail, and mucosal findings in children with Turner syndrome (TS) and their associations with coexisting endocrinopathies.

Methods: A cross-sectional study was conducted with TS patients who were followed up by pediatric endocrinology and referred to the dermatology outpatient clinic. Data were collected using standardized dermatological examination forms, including demographic information, clinical features, presence of endocrinopathies, and medication usage. Statistical analysis was used to evaluate differences between groups and to analyze relationships between variables.

Results: The cohort consisted of 112 girls with TS. Skin and hair findings were detected in 86.6% of the patients, with melanocytic nevi (44.6%) and xerosis (41.1%) being the most commonly observed. Oral mucosal findings were observed less frequently (17.0%). Nail findings were detected in 63.4% of the cases, with leukonychia (15.2%) and subungual hyperkeratosis (14.3%) being the most prevalent. Older age, delayed diagnosis, longer follow-up duration, and lower body mass index were associated with an increased frequency of skin and hair findings ($p<0.05$). Moreover, the presence of coexisting endocrinopathies was significantly associated with skin and hair findings. Nail findings were significantly associated with longer follow-up duration ($p=0.002$), the presence of endocrinopathies ($p<0.001$), and comorbidities ($p=0.004$).

Conclusion: This study confirmed that skin, hair, and nail abnormalities are commonly observed in TS. The association of these findings with endocrinopathies suggests that systemic factors influence dermatological problems in TS. It is recommended to integrate dermatological evaluations into the routine endocrine and cardiological follow-up of children with TS.

Keywords: Turner syndrome, skin findings, endocrinopathies, melanocytic nevus, nail findings

What is already known on this topic?

Turner syndrome (TS) is characterized by a constellation of morphological clinical features, including short stature, webbed neck, low posterior hairline, and widely spaced nipples. Existing literature documents associations between TS and increased melanocytic nevi, halo nevi, ichthyotic changes, pilomatricoma, café-au-lait spots, and hypertrichosis. Endocrinopathies are common in Turner syndrome. The relationship between these endocrinopathies and cutaneous findings is not clear in the literature.

What this study adds?

Oral mucosal findings were present in 17% of TS children. The most frequent findings were geographic tongue, fungiform papilla hypertrophy, atrophic glossitis, and scrotal tongue. There are no known studies specifically examining oral mucosal findings in TS patients, making our study the only one on the subject. In this study, nail findings were seen in 63.4%, with leukonychia most frequently observed, followed by subungual hyperkeratosis and pincer nail, a characteristic TS finding. The significant association between skin and hair findings and concomitant endocrinopathies ($p=0.001$) observed in this study supports the hypothesis of an association.

Introduction

Turner syndrome (TS) is a genetic disorder that affects females and is caused by the complete or partial loss of one X chromosome and is one of the most common chromosomal anomalies (1). The syndrome is characterized by a range of morphological clinical features (stigmata), including short stature, webbed neck, low posterior hairline, and widely spaced nipples (1). Affected individuals frequently present with linear growth retardation, gonadal dysgenesis, hypothyroidism, osteoporosis, and skeletal, cardiovascular, and renal anomalies. Early-onset sensorineural hearing loss, as well as neurocognitive and psychosocial issues, are also common systemic manifestations (2).

Although skin, hair, nail, and mucosal changes are commonly seen in TS, these findings are often under-recognized during patient follow-up. In the largest multicenter cohort reported from

Turkey, dermatological problems were reported in 21.8% of cases (3). Existing literature has documented associations between TS and increased melanocytic nevi, halo nevi, ichthyosiform changes, pilomatricomas, café-au-lait spots, and hypertrichosis (4,5,6,7). However, a comprehensive understanding of the relationship between these dermatological findings and coexisting endocrinopathies remains limited. Moreover, the increased predisposition to autoimmunity in TS may elevate the risk of conditions such as vitiligo and alopecia areata (8,9). Nevertheless, strong evidence-based data supporting these associations are lacking.

The aim of this study was to comprehensively evaluate skin, hair, nail, and mucosal findings in children and adolescents with TS followed in a pediatric endocrinology clinic and to investigate the potential associations between these findings and coexisting endocrinopathies. Enhancing the knowledge

base regarding these findings may contribute to improving diagnostic and therapeutic approaches for children with TS.

Materials and Methods

This study involved a cross-sectional assessment of cutaneous, hair, nail, and mucosal findings in girls diagnosed with TS, all of whom were followed by a pediatric endocrinologist and subsequently referred to the dermatology outpatient clinic. Findings were collected from nine tertiary care centres. Exclusion criteria included: age over 18 years, male sex, pregnancy, lactation, prior epilation, and use of oral contraceptives, corticosteroids, cyclosporine, or spironolactone within the preceding three months.

All participants underwent thorough dermatological evaluations performed by a senior dermatologist, using standardized examination forms and routine dermatological examination techniques. Pertinent clinical data, including age, age at diagnosis, follow-up duration, medication history, height, weight, body mass index (BMI), comorbidities, and endocrinopathies (diabetes, hyperlipidemia, hypertension, thyroid disease), were systematically recorded. Standardized methods were consistently employed for dermatological evaluations across all participants. Informed consent was obtained from all participants and their parents prior to enrollment. Ethical approval was granted by the Necmettin Erbakan University Non-Drug and Non-Medical Device Research Ethics Committee (approval no.: 2021/3023, date: 22.01.2021).

Statistical Analysis

Data analysis was performed using SPSS (Statistical Package for the Social Sciences), version 18.0 (SPSS Inc., Chicago, ILL, USA). Statistical methods were employed to assess intergroup differences and analyze relationships between variables. Descriptive statistics included frequencies (n) and percentages (%) for categorical variables, and means±standard deviations (SD) and interquartile ranges (1st-3rd quartile) for numerical variables. Categorical variable distributions were examined using Pearson's chi-square test and Fisher's exact test. For non-normally distributed numerical variables, comparisons between two independent groups were performed using the Mann-Whitney U test, and comparisons among more than two independent groups were conducted using the Kruskal-Wallis test. In cases where the Kruskal-Wallis test yielded significant results, post-hoc analysis was performed using the Mann-Whitney U test with Dunn-Bonferroni correction. A p-value of <0.05 was considered indicative of statistical significance for all analyses.

Results

This study encompassed 112 children and adolescents diagnosed with Turner syndrome. Comprehensive examinations

were conducted to assess skin, hair, oral mucosa, and nail characteristics. Table 1 presents the distribution of demographic and clinical characteristics. Karyotype analysis revealed the presence of 45,X in 48.2% (n=54) of cases, mosaic X chromosome in 32.1% (n=36), and X chromosome anomalies in 7.1% (n=8). Comorbidities were evident in 39.3% (n=44) of the Turner syndrome cohort, with cardiopathy representing the most common comorbidity (15.2%, n=17). Growth hormone (GH) therapy constituted the predominant treatment modality, administered to 50.0% (n=56) of the patients.

Dermatological findings are summarized in Table 2. Skin and hair abnormalities were identified in 86.6% (n=97) of the cohort, with melanocytic nevus (44.6%, n=50), xerosis (41.1%, n=46), and keratosis pilaris (25.9%, n=29) representing the most

Table 1. Demographic and clinical characteristics of Turner syndrome cases

Variables		Results (n=112)
Age (years); (mean±SD; IQR)		13.33±4.60 (11-18)
Sex; n (%)	Female	112 (100.0)
Karyotype; n (%)	45,X	54 (48.2)
	Mosaicism	36 (32.1)
	X chromosome anomaly	8 (7.1)
	Unknown	14 (12.5)
BMI; (mean±SD)		23.11±18.90
Age at diagnosis (years); (mean±SD; IQR)		6.93±5.01 (1-11)
Follow-up duration (months); (mean±SD; IQR)		79.17±64.61 (24-120)
Comorbidities; n(%)		44 (39.3)
Cardiopathy		17 (15.2)
Obesity		7 (6.3)
Kidney anomaly		7 (6.3)
Mental retardation		7 (6.3)
Celiac disease		5 (4.5)
Hyperlipidemia		3 (2.7)
Selective IgM deficiency		2 (1.8)
Allergic rhinitis		2 (1.8)
FMF		1 (0.9)
Hypertension		1 (0.9)
ALL		1 (0.9)
Endocrinopathies; n (%)		74 (66.1)
GH deficiency		40 (35.7)
Hyperthyroidism		26 (23.2)
Hypothyroidism		16 (14.3)
Diabetes		6 (5.4)
Osteoporosis		2 (1.8)
Medication use ^(a)		
Growth hormone therapy n (%)		56 (50.0)
Estrogen n (%)		25 (22.3)
Thyroid medication n (%)		15 (13.4)
Other n (%)		13 (11.6)

Footnote: a) The medication categories are not mutually exclusive; some cases involve the use of more than one medication.

BMI: body mass index, FMF: Familial Mediterranean Fever, ALL: acute lymphocytic leukemia, GH: growth hormone, IQR: interquartile range

frequent findings. Oral mucosal findings were detected in 17.0% (n=19) of participants, with geographic tongue and fungiform papilla hypertrophy each observed in 3.6% (n=4). Nail findings were observed in 63.4% (n=71) of all cases, the most common ones were, leukonychia (15.2%, n=17), subungual hyperkeratosis (14.3%, n=16), and pincer nail (10.7%, n=12).

Table 3 shows the distribution of skin, hair, and oral mucosal findings in relation to demographic and clinical characteristics.

The prevalence of skin and hair abnormalities exhibited statistically significant elevations in cases characterized by older age, delayed diagnosis, longer follow-up duration, and lower BMI ($p<0.001$, $p=0.001$, $p=0.002$, $p=0.049$, respectively). Moreover, cases with concomitant endocrinopathy demonstrated significantly increased prevalence of skin and hair abnormalities ($p=0.001$). Older age and lower BMI were statistically associated with the prevalence of oral mucosal findings ($p=0.002$, $p=0.009$, respectively).

Table 2. Distribution of dermatological findings in patients

Variables	Results (n=112)	Variables	Results (n=112)
Skin and hair findings; n (%)	97 (86.6)		
Melanocytic nevus	50 (44.6)		
Xerosis	46 (41.1)		
Keratosis pilaris	29 (25.9)		
Seborrheic dermatitis	27 (24.1)		
Pruritus	21 (18.8)		
Striae	18 (16.1)		
Acne	15 (13.4)		
Telogen effluvium	9 (8.0)		
Dysplastic nevus	7 (6.3)		
Atopic dermatitis	7 (6.3)		
Hypopigmentation	7 (6.3)		
Café-au-lait spots	7 (6.3)		
Psoriasis	6 (5.4)		
Hirsutism	5 (4.5)		
Pityriasis alba	5 (4.5)		
Pityriasis versicolor	5 (4.5)		
Traction alopecia	4 (3.6)		
Urticaria	4 (3.6)		
Eczema	3 (2.7)		
Congenital nevus	3 (2.7)		
Hemangioma	3 (2.7)		
Alopecia areata	3 (2.7)		
Androgenetic alopecia	3 (2.7)		
Hyperpigmentation	3 (2.7)		
Lymphedema	3 (2.7)		
Acanthosis nigricans	2 (1.8)		
Verruca (Wart)	2 (1.8)		
Halo nevus	1 (0.9)		
Skin tag	1 (0.9)		
Trichotillomania	1 (0.9)		
Herpes labialis	1 (0.9)		
Tinea pedis	1 (0.9)		
Keloid	1 (0.9)		
Other	11 (9.8)		
		Nail findings; n (%)	71 (63.4)
		Leukonychia	17 (15.2)
		Subungual hyperkeratosis	16 (14.3)
		Pincer nail	12 (10.7)
		Unguis incarnatus	9 (8.0)
		Longitudinal ridging	8 (7.1)
		Terry's nails	8 (7.1)
		Brittle nails	8 (7.1)
		Transverseridging	7 (6.3)
		onycholysis	6 (5.4)
		Pitting	5 (4.5)
		Increased nail curvature	4 (3.6)
		Trachyonychia	3 (2.7)
		Onychophagia	2 (1.8)
		Onychomadesis	1 (0.9)
		Beau's lines	1 (0.9)
		Koilonychia (Spoon nails)	1 (0.9)
		Muehrcke's lines	1 (0.9)
		Mees' lines	1 (0.9)
		Lindsay's nails	1 (0.9)
		Congenital malalignment	1 (0.9)
		Racquet nail	1 (0.9)
		Other	4 (3.6)
Oral mucosal findings; n (%)	19 (17.0)	Affected nail location; n (%)	
Geographic tongue	4 (3.6)	Hand	12 (10.7)
Fungiform papilla hypertrophy	4 (3.6)	Foot	29 (25.9)
Atrophic glossitis	3 (2.7)	Both hand and foot	30 (26.8)
Scrotal tongue	3 (2.7)		
Candidiasis	2 (1.8)		
Morsicatio buccarum	2 (1.8)		
Gingival hypertrophy	2 (1.8)		
Aphthous ulcer	2 (1.8)		
Angular cheilitis	1 (0.9)		
Exfoliative cheilitis	1 (0.9)		
Gingival bleeding	1 (0.9)		
Lichen planus	1 (0.9)		
		Number of affected nails; (mean±SD; IQR)	6.81±6.75 (0.7-10.0)

SD: standard deviation, IQR: interquartile range

Table 4 presents the distribution of nail findings in relation to demographic and clinical characteristics. Nail finding prevalence was significantly more common in patients with longer follow-up duration ($p=0.002$) and a higher prevalence of associated comorbidities and endocrinopathies ($p=0.004$, $p<0.001$, respectively). A significant difference was found in age distribution according to the affected nail location ($p=0.044$). Participants with hand nail findings only were younger than those with both hand and foot nail findings ($p<0.05$). In girls using medications, the prevalence of foot nail findings was significantly more common ($p=0.014$).

Discussion

This study evaluated the frequency of mucocutaneous findings in children with TS and associations with demographic and clinical variables. Skin and hair findings were detected in more than four fifths, while nail findings were observed in approximately two-thirds of patients. Oral mucosal findings were observed at a much lower frequency (17.0%). Our findings suggest that the mucocutaneous spectrum in this TS cohort was broad, and skin-hair and nail findings should be considered together with accompanying clinical features. The detrimental effects of endocrine disorders on the skin, hair, and mucosal

Table 3. Distribution of skin, hair, and oral mucosal findings according to demographic and clinical characteristics

	Skin & hair findings			Oral mucosal findings		
	Present	Absent	p value	Present	Absent	p value
Age (years); (mean±SD; IQR)	14.36±4.71 (12-17)	6.67±6.45 (1-12)	<0.001*	16.42±4.61 (15-20)	12.70±5.59 (10-17)	0.002*
Karyotype; n (%)						
45,X	42 (77.8)	12 (22.2)	0.096**	9 (16.7)	45 (83.3)	0.455**
Mosaicism	34 (94.4)	2 (5.6)		6 (16.7)	30 (83.3)	
X chromosome anomaly	7 (87.5)	1 (12.5)		0 (0.0)	8 (100.0)	
BMI; (mean±SD; IQR)	21.05±4.47 (17.9-23.5)	36.38±49.78 (15.1-23.6)	0.049*	22.85±4.06 (20.4-25.8)	23.16±20.68 (17.1-22.9)	0.009*
Age at diagnosis (years); (mean±SD; IQR)	7.53±4.91 (3-12)	2.86±3.65 (0-5)	0.001*	7.53±5.94 (1-12.5)	6.82±4.85 (2.5-11.0)	0.285*
Follow-up duration (months); (mean±SD; IQR)	84.99±64.15 (36-120)	39.29±54.41 (12-39)	0.002*	90.06±62.28 (42-144)	77.18±65.15 (24-120)	0.407*
Associated comorbidity; n (%)	39 (88.6)	5 (11.4)	0.612**	10 (22.7)	34 (77.3)	0.191**
Associated endocrinopathy; n (%)	70 (94.6)	4 (5.4)	0.001**	15 (20.3)	59 (79.7)	0.193**
Medication use; n (%)	76 (89.4)	9 (10.6)	0.190***	16 (18.8)	69 (81.2)	0.556***

n (%); mean±SD; IQR, *Mann-Whitney U test, **Pearson chi-square test, ***Kruskal-Wallis test
SD: standard deviation, IQR: interquartile range

Table 4. Distribution of nail findings according to demographic and clinical characteristics

	Nail findings			Affected nail location			
	Present	Absent	p value	Hand	Foot	Hand+Foot	p value
Age (years); (mean±SD; IQR)	14.27±5.32 (11-18)	11.71±5.75 (8-16)	0.068*	12.58±3.26 (11-15)	13.38±4.98 (10-17)	15.80±6.00 (12-20)	0.044***
BMI; (mean±SD; IQR)	21.27±4.38 (17.9-23.8)	26.28±30.69 (16.5-21.9)	0.064*	21.78±6.35 (17.2-23.1)	20.83±4.45 (17.2-23.9)	21.49±3.39 (18.7-24.1)	0.641***
Age at diagnosis (years); (mean±SD; IQR)	6.44±4.69 (1.5-7.0)	7.76±5.46 (1.0-12.5)	0.197*	8.00±3.43 (5-11)	5.90±4.83 (0.4-9)	6.37±4.95 (1-11)	0.454***
Follow-up duration (months); (mean±SD; IQR)	92.71±67.36 (38-150)	56.39±53.03 (15-78)	0.002*	57.09±40.71 (36-72)	87.32±67.96 (24-150)	110.80±70.12 (60-162)	0.064***
Comorbidity; n (%)	35 (79.5)	9 (20.5)	0.004**	4 (11.4)	18 (51.4)	13 (37.1)	0.170**
Endocrinopathy; n (%)	57 (77.0)	17 (23.0)	<0.001**	9 (15.8)	24 (42.1)	24 (42.1)	0.850**
Medication use; n (%)	56 (65.9)	29 (34.1)	0.332**	12 (21.4)	25 (44.6)	19 (33.9)	0.014**

n (%); mean±SD; IQR, *Mann-Whitney U test, **Pearson chi-square test, ***Kruskal-Wallis test
SD: standard deviation, IQR: interquartile range

tissues are well known (10,11). In Turner syndrome, frequently encountered thyroid disorders and growth retardation may also be more commonly associated with mucocutaneous findings (6,8). Indeed, in our study, the significantly higher frequency of skin–hair findings associated with the presence of an endocrinopathy ($p=0.001$) supports this observation.

While skin–hair findings were associated with older age, diagnostic delay, and presence of endocrinopathy; oral mucosal findings were more closely associated with older age and low BMI. Nail findings significantly increased, particularly in the presence of endocrinopathy and comorbidities, suggesting that nail changes may be one of the most sensitive dermatological indicators of systemic involvement in Turner syndrome.

Prevalence data regarding dermatological findings in Turner syndrome vary widely due to methodological differences. Studies based on medical record reviews report lower prevalence rates, while dermatology-focused studies using detailed physical examination or patient/parent reporting have shown rates of findings such as xerosis and melanocytic nevi reaching 70–80% (3,8,12). This discrepancy may be explained because mild or subclinical findings are not always recorded as “clinical problems” during routine follow-up visits. In the present study, the use of a standardized dermatological examination form by a dermatologist likely contributed to the higher detection rates. Furthermore, the differences may also be affected by referral bias and inter-center variability in assessment practices.

The most common skin finding in our study cohort was melanocytic nevi, observed in about half of the patients. Benign acquired nevi were most frequently seen, while dysplastic and congenital nevi were less common. Although previous reports have indicated an increased prevalence of halo nevi in Turner syndrome, this was found in only one case in our study (9). Earlier studies have shown that the average number of nevi is higher in individuals with TS compared to the general population (3,8). Considering that the total number of melanocytic nevi is an independent risk factor for melanoma, it would be appropriate to include regular nevus monitoring in the follow-up of individuals with TS. However, there is currently no evidence suggesting a significantly increased risk of melanoma in girls and women with TS (4,13,14,15).

Growth hormone (GH) therapy is a commonly used approach in the management of short stature in TS and was also the most frequently applied treatment in our study. Some reports have suggested that GH treatment may increase the number of nevi in TS (16,17,18). However, the causal nature of this relationship remains unclear and further studies are needed to determine whether this is due to growth axis abnormalities or the therapy itself.

The frequent occurrence of xerosis cutis and keratosis pilaris among the most common dermatological findings suggests that keratinization processes may be widely affected in TS. As Borroni et al. (8) reported, skin dryness in TS should not only be considered as non-specific xerosis but also as a manifestation of ichthyosis-like keratinization disorders. In a more recent study, 78.7% of individuals with TS reported complaints of dry, scaly, or flaky skin, with this being one of the factors most affecting quality of life (12). Nonetheless, the pathophysiological relationship between xerosis and TS has not yet been fully elucidated.

A large-scale cohort study conducted in Denmark showed that inflammatory skin diseases, such as psoriasis, atopic dermatitis, and seborrheic dermatitis, were reported at higher rates in individuals with TS compared to the general population (19). In the present study, the relatively high frequency of seborrheic dermatitis (24.1%) may be considered consistent with this literature, suggesting that GH and the insulin-like growth factor (IGF) axis affect sebaceous gland function and sebum production (20). However we were not able to test whether TS patients who received GH treatment had higher frequency of these problems than those who did not receive GH treatment.

The detection of acne vulgaris in 13.4% of cases in our study suggests that this finding occurs at a relatively low frequency in TS, consistent with existing literature (21). This may be related to reduced sebum production, low peripheral androgen levels, and the suppression of testosterone and dihydrotestosterone levels due to hormone replacement therapy in TS patients (21,22).

Lymphedema is frequently observed at birth in Turner syndrome and tends to regress in most cases by the age of two. Therefore, its low detection rate (2.7%) in our study was an expected finding.

Among hair-related dermatological findings, telogen effluvium was the most common (8.0%). Hormonal imbalances, such as estrogen deficiency, thyroid dysfunction, and abnormalities in the growth axis may affect the hair follicle cycle and increase transition to the telogen phase. This finding indicates that complaints related to hair should be systematically questioned during dermatological evaluations in TS patients.

Oral mucosal findings were observed in 17.0% of TS cases in our study. The most common findings included geographic tongue, hypertrophy of fungiform papillae, atrophic glossitis, and fissured tongue. Data evaluating oral mucosal findings in a systematic manner in TS are limited; therefore, our findings contribute to the existing evidence of this association. The positive association of oral mucosal findings with age and their negative association with BMI are noteworthy. This relationship may be linked to secondary nutritional deficiencies and micronutrient deficits. Particularly, findings such as atrophic glossitis and angular cheilitis are classic indicators of micronutrient deficiencies.

However, due to the lack of biochemical assessments in our study, further investigations are needed to confirm these associations.

Nail findings were observed in 63.4% of children with Turner syndrome in our study. The most frequent nail finding was leukonychia, followed by subungual hyperkeratosis and pincer nails, which are considered characteristic of TS. The significantly higher frequency of nail findings in the presence of endocrinopathy and their strong association with comorbidities suggest that nail changes may be a sensitive marker of systemic disease burden in Turner syndrome.

Circulatory disorders, hormonal imbalances, and connective tissue anomalies commonly reported in TS may negatively affect keratin production and nail structure. Hypoplastic and dysplastic nails have also been reported to be associated with lymphedema in the literature (23). Moreover, the wide and short foot structure common in TS, lymphedema, and hyperextension of the big toe may predispose individuals to mechanical trauma due to increased shoe pressure, potentially contributing to the development of leukonychia, subungual hyperkeratosis, and pincer nails (24).

In the present study, the observation that nail findings were more frequent in cases with a longer follow-up duration, and that patients with only fingernail involvement were younger than those with both hand and foot involvement, suggests that nail findings may be related to cumulative systemic and mechanical effects over time. In light of these findings, we believe that nail examination should not be neglected in the dermatological evaluation of individuals with Turner syndrome and should be considered a potential marker of systemic involvement.

Study Limitations

The main limitation of this study is the lack of a control group, which prevents direct comparison of the observed mucocutaneous findings with their frequencies in the general population. In addition, the cross-sectional design does not allow for causal inference between dermatological findings and endocrinopathies.

Endocrinopathies and comorbidities were recorded based on clinical diagnoses established by pediatric endocrinology. The presence of thyroid autoantibodies, hormone levels, IGF-1 measurements, and/or dynamic endocrine tests were not analyzed within the scope of this study. Thus, findings such as the higher reported frequency of hyperthyroidism compared to hypothyroidism may reflect differences in clinical records or classification rather than true prevalence. Considering that short stature in TS is often due to functional impairment of the growth axis rather than biochemical GH deficiency, some cases recorded as “GH deficiency” may instead reflect growth failure.

Endocrinopathies were not analyzed according to subtypes. The presence of multiple endocrine disorders in many cases may have introduced confounding effects. The complexity resulting from this and the lack of sufficient statistical power for subgroup analyses limited the ability to isolate specific associations between dermatological findings and individual endocrine conditions. Lastly, the absence of biochemical evaluations related to nutritional status and possible micronutrient deficiencies, such as iron, B-complex vitamins, and zinc, limited the interpretation of the association between low BMI and oral mucosal findings.

Conclusion

Endocrinopathies in individuals with TS were associated with an increase in the frequency of dermatological findings, particularly those involving the skin, hair, and nails. Therefore, in addition to routine endocrinology and cardiology follow-ups, regular dermatological evaluations may be appropriate for the optimal care of girls, adolescents and women with TS.

Ethics

Ethics Committee Approval: Ethical approval was granted by the Necmettin Erbakan University Non-Drug and Non-Medical Device Research Ethics Committee (approval no.: 2021/3023, date: 22.01.2021).

Informed Consent: Informed consent was obtained from all participants and their parents prior to enrollment.

Footnotes

Authorship Contributions

Surgical and Medical Practices: Selami Aykut Temiz, Filiz Cebeci, Aslıhan Çiçekli, Concept: Selami Aykut Temiz, Filiz Cebeci, Aslıhan Çiçekli, Design: Selami Aykut Temiz, Filiz Cebeci, Aslıhan Çiçekli, Begüm Işık, Sevil Savaş Erdoğan, Selma Emre, Hilal Kaya Erdoğan, İsa An, Sezgi Sarıkaya Solak, Sevilay Kılıç, Eda Öksüm Solak, Arzu Ataseven, Beray Selver Eklioğlu, Suna Kılınç, Mehmet Boyraz, Enver Şimşek, Tuğba Çetin Kontbay, Filiz Tütüncüler, Durmuş Doğan, Ülkü Gül Şiraz, Güllü Eren, Data Collection or Processing: Selami Aykut Temiz, Filiz Cebeci, Aslıhan Çiçekli, Begüm Işık, Sevil Savaş Erdoğan, Selma Emre, Hilal Kaya Erdoğan, İsa An, Sezgi Sarıkaya Solak, Sevilay Kılıç, Eda Öksüm Solak, Arzu Ataseven, Beray Selver Eklioğlu, Suna Kılınç, Mehmet Boyraz, Enver Şimşek, Tuğba Çetin Kontbay, Filiz Tütüncüler, Durmuş Doğan, Ülkü Gül Şiraz, Güllü Eren, Analysis or Interpretation: Selami Aykut Temiz, Filiz Cebeci, Aslıhan Çiçekli, Begüm Işık, Sevil Savaş Erdoğan, Selma Emre, Hilal Kaya Erdoğan, İsa An, Sezgi Sarıkaya Solak, Sevilay Kılıç, Eda Öksüm Solak, Arzu Ataseven, Beray Selver Eklioğlu, Suna Kılınç, Mehmet Boyraz, Enver Şimşek, Tuğba Çetin Kontbay, Filiz Tütüncüler, Durmuş Doğan, Ülkü Gül Şiraz, Güllü Eren, Literature Search: Selami Aykut Temiz, Filiz Cebeci, Aslıhan Çiçekli, Writing: Selami Aykut Temiz, Filiz Cebeci, Aslıhan Çiçekli.

Financial Disclosure: The authors declare that the work reported in this manuscript was undertaken with their financial resources, and they did not receive any financial help from any individual, group, or agency.

References

1. Ranke MB, Saenger P. Turner's syndrome. *Lancet*. 2001;358:309-314.
2. Gravholt CH, Andersen NH, Conway GS, Dekkers OM, Geffner ME, Klein KO, Lin AE, Mauras N, Quigley CA, Rubin K, Sandberg DE, Sas TCJ, Silberbach M, Söderström-Anttila V, Stochholm K, van Alfen-van derVelden JA, Woelfle J, Backeljauw PF; International Turner Syndrome Consensus Group. Clinical practice guidelines for the care of girls and women with Turner syndrome: proceedings from the 2016 Cincinnati International Turner Syndrome Meeting. *Eur J Endocrinol*. 2017;177:G1-G70.
3. Y Yeşilkaya E, Bereket A, Darendeliler F, Baş F, Poyrazoğlu Ş, Küçükemre Aydın B, Darcan Ş, Dünder B, Büyükinan M, Kara C, Sarı E, Adal E, Akıncı A, Atabek ME, Demirel F, Çelik N, Özkan B, Özhan B, Orbak Z, Ersoy B, Doğan M, Ataş A, Turan S, Gökşen D, Tarım Ö, Yüksel B, Ercan O, Hatun Ş, Şimşek E, Ökten A, Abacı A, Döneray H, Özbek MN, Keskin M, Önal H, Akyürek N, Bulan K, Tepe D, Emeksiz HC, Demir K, Kızılay D, Topaloğlu AK, Eren E, Özen S, Abalı S, Akın L, Selver Eklioğlu B, Kaba S, Anık A, Baş S, Ünüvar T, Sağlam H, Bolu S, Özgen T, Doğan D, Deniz Çakır E, Şen Y, Andıran N, Çizmecioglu F, Evliyaoğlu O, Karagüzel G, Pirgon Ö, Çatlı G, Can HD, Gürbüz F, Binay Ç, Baş VN, Fidancı K, Polat A, Gül D, Açıkel C, Demirbilek H, Cınaz P, Bondy C. Turner syndrome and associated problems in Turkish children: a multicenter study. *J Clin Res Pediatr Endocrinol*. 2015;7:27-36.
4. Becker B, Jospe N, Goldsmith LA. Melanocytic nevi in Turner syndrome. *Pediatr Dermatol*. 1994;11:120-124.
5. Gunther DF, Sybert VP. Lymphatic, tooth and skin manifestations in Turner syndrome. *International Congress Series*, Elsevier, 2006.
6. Lowenstein EJ, Kim KH, Glick SA. Turner's syndrome in dermatology. *J Am Acad Dermatol*. 2004;50:767-776.
7. Wood S, Nguyen D, Hutton K, Dickson W. Pilomatricomas in Turner syndrome. *Pediatr Dermatol*. 2008;25:449-451.
8. Borroni G, Larizza D, Vignati G, Zaccane C, Zanetta R, Vignoli GP, Brazzelli V, Bobba C, Rabbiosi G. The dermatological spectrum of Turner syndrome. *J Eur Acad Dermatol Venereol*. 1994;3:334-340.
9. Brazzelli V, Larizza D, Martinetti M, Martinoli S, Calcaterra V, De Silvestri A, Pandolfi R, Borroni G. Halo nevus, rather than vitiligo, is a typical dermatologic finding of turner's syndrome: clinical, genetic, and immunogenetic study in 72 patients. *J Am Acad Dermatol*. 2004;51:354-358.
10. Jabbour SA. Cutaneous manifestations of endocrine disorders: a guide for dermatologists. *Am J Clin Dermatol*. 2003;4:315-331.
11. Lause M, Kamboj A, Fernandez Faith E. Dermatologic manifestations of endocrine disorders. *Transl Pediatr*. 2017;6:300-312.
12. Rodriguez-Buritica D, Mones M, Prakash SK, Rivera M, Aldrich M, Rogge M, Richardson K. Dermatological concerns for women and girls with turner syndrome. *Front Med (Lausanne)*. 2023;10:1235187.
13. Holly EA, Kelly JW, Shpall SN, Chiu SH. Number of melanocytic nevi as a major risk factor for malignant melanoma. *J Am Acad Dermatol*. 1987;17:459-468.
14. Bauer J, Garbe C. Acquired melanocytic nevi as risk factor for melanoma development. A comprehensive review of epidemiological data. *Pigment Cell Res*. 2003;16:297-306.
15. Viuff MH, Stochholm K, Lin A, Berglund A, Juul S, Gravholt CH. Cancer occurrence in Turner syndrome and the effect of sex hormone substitution therapy. *Eur J Endocrinol*. 2021;184:79-88.
16. Ali I, Moore T, Kashyap A, Starling C, Thomas J, Truong VTT, Prakash SK, Rogge MN. Increased acquired melanocytic nevi in Turner syndrome associated with use of growth hormone: An observational study. *J Am Acad Dermatol*. 2024;91:744-746. Epub 2024 Jun 19
17. Piérard GE, Piérard-Franchimont C, Nikkels A, Nikkels-Tassoudji N, Arrese JE, Bourguignon JP. Naevocyte triggering by recombinant human growth hormone. *J Pathol*. 1996;180:74-79.
18. Wyatt D. Melanocytic nevi in children treated with growth hormone. *Pediatrics*. 1999;104(Suppl 5):1045-1050.
19. Viuff MH, Stochholm K, Juul S, Gravholt CH. Disorders of the eye, ear, skin, and nervous system in women with Turner syndrome -a nationwide cohort study. *Eur J Hum Genet*. 2022;30:229-236. Epub 2021 Oct 28
20. Deplewski D, Rosenfield RL. Growth hormone and insulin-like growth factors have different effects on sebaceous cell growth and differentiation. *Endocrinology*. 1999;140:4089-4094.
21. Brazzelli V, Larizza D, Muzio F, Calcaterra V, Fornara L, Klersy C, Borroni G. Low frequency of acne vulgaris in adolescent girls and women with Turner's syndrome: a clinical, genetic and hormonal study of 65 patients. *Br J Dermatol*. 2008;159:1209-1211. Epub 2008 Sep 15
22. Brazzelli V, Calcaterra V, Muzio F, Klersy C, Larizza D, Borroni G. Reduced sebum production in Turner syndrome: a study of twenty-two patients. *Int J Immunopathol Pharmacol*. 2011;24:789-792.
23. Ikomi C, Blatt J, Ghofrani S, Zhang R, Ross J, Law JR. Lymphedema in Turner syndrome: correlations with phenotype and karyotype. *J Pediatr Endocrinol Metab*. 2024;37:789-795.
24. Findlay CA, Donaldson MD, Watt G. Foot problems in Turner's syndrome. *J Pediatr*. 2001;138:775-777.