



DOI: 10.4274/jcrpe.galenos.2026.2026-4-6

J Clin Res Pediatr Endocrinol 2026;18(3):554-555

Anorexia Nervosa Emerging after Swyer Syndrome Diagnosis

Demet Aygün Arı¹, Semra Çetinkaya²

¹Hacettepe University Faculty of Medicine, Department of Pediatrics, Division of Adolescent Medicine, Ankara, Türkiye

²University of Health Sciences Türkiye, Ankara Etlik City Hospital, Clinic of Pediatrics, Division of Pediatric Endocrinology, Ankara, Türkiye

Cite this article as: Aygün Arı D, Çetinkaya D. Anorexia nervosa emerging after Swyer syndrome diagnosis. J Clin Res Pediatr Endocrinol. [Epub Ahead of Print] 2026;18(3):554-555

Keywords: Swyer syndrome, complete gonadal dysgenesis, anorexia nervosa, psychiatric comorbidity

Dear Editor,

Swyer syndrome (46,XY complete gonadal dysgenesis) is a rare disorder of sex development (DSD) characterized by female external genitalia, hypergonadotropic hypogonadism, primary amenorrhea, and increased risk of gonadal malignancy (1,2). Beyond endocrine and oncologic complications, psychosocial difficulties are increasingly recognized in individuals with DSD. We report an adolescent with Swyer syndrome who developed anorexia nervosa during follow-up after diagnosis and treatment.

A 15.5-year-old phenotypic female presented with primary amenorrhea despite spontaneous breast and pubic hair development since age 11 years. Physical examination showed Tanner stage IV breast and pubic hair development with normal female external genitalia. Laboratory evaluation demonstrated hypergonadotropic hypogonadism (follicle-stimulating hormone 138.07 mIU/mL, luteinizing hormone 48.35 mIU/mL, estradiol <12.1 pg/mL). Pelvic imaging revealed absent ovarian tissue and a left adnexal nodular lesion. Karyotype analysis showed 46,XY with positive SRY expression. During the diagnostic process, tumor markers increased (alpha-fetoprotein 5.6 ng/mL, β -hCG 6.6 mIU/mL), raising suspicion for gonadal malignancy.

Laparoscopic bilateral gonadectomy was performed, and histopathological examination demonstrated gonadoblastoma in both gonads with coexisting dysgerminoma in the left gonad. Postoperatively, estrogen replacement therapy was initiated and later transitioned to cyclic estrogen-progesterone therapy.

At initial psychosocial assessment, no apparent psychiatric risk factors were identified. However, six months after diagnosis, she had intentionally lost 16 kg over three months through severe caloric restriction. At that time, her weight was 45 kg [-1.94 standard deviation score (SDS); 2.6th percentile], height was 164 cm (0.26 SDS; 60.3rd percentile), and body mass index (BMI) was 16.7 kg/m² (-2.63 SDS; 0.4th percentile), corresponding to 78.8% of median BMI. Further psychosocial assessment revealed increasing distress related to her diagnosis, concerns regarding gender identity, and dissatisfaction with physical appearance. She demonstrated distorted body image and intense fear of gaining weight and was diagnosed with anorexia nervosa according to the fifth revision of the Diagnostic and Statistical Manual of Mental Disorders criteria. As she remained medically stable, outpatient management was initiated. Her caloric intake was gradually increased, and she was closely monitored during

Corresponding Author: Demet Aygün Arı, MD, PhD, Hacettepe University Faculty of Medicine, Department of Pediatrics, Division of Adolescent Medicine, Ankara, Türkiye

E-mail: demetaygunari@gmail.com **ORCID:** orcid.org/0000-0002-8701-6085

Conflict of interest: None declared

Received: 11.04.2026 **Accepted:** 16.05.2026 **Epub:** 22.05.2026 **Publication Date:** 08.09.2026



©Copyright 2026 by Turkish Society for Pediatric Endocrinology and Diabetes / The Journal of Clinical Research in Pediatric Endocrinology published by Galenos Publishing House. Licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 (CC BY-NC-ND) International License.

follow-up. No signs of refeeding syndrome or other medical complications were observed during nutritional rehabilitation. She was also evaluated regularly by psychiatry and dietetics teams as part of multidisciplinary care. During follow-up, she regained a healthy weight and maintained positive body image.

Patients with DSD are known to experience increased rates of psychological distress, including anxiety, depression, stigma, impaired self-esteem, and body dissatisfaction (3,4,5). Current consensus guidelines emphasize psychosocial support as part of multidisciplinary care (4). A large European study reported eating disorders in 11.1% of adults with DSD, particularly among individuals with an XY karyotype raised as females (3). In contrast, studies in adolescents with DSD have focused mainly on anxiety, depressive symptoms, and attention-deficit/hyperactivity disorder, while eating disorders have rarely been discussed (5). To the best of our knowledge, reports of anorexia nervosa in adolescents with Swyer syndrome are scarce (6). Our case highlights that eating disorders may emerge shortly after diagnosis and treatment, even in adolescents without previous psychosocial risk factors.

This report highlights the importance of multidisciplinary management in Swyer syndrome, including routine psychosocial

assessment during follow-up. Clinicians caring for adolescents with DSD should remain aware of the potential for eating disorders and body image concerns during the diagnostic and treatment process.

References

1. Michala L, Goswami D, Creighton SM, Conway GS. Swyer syndrome: presentation and outcomes. *BJOG*. 2008;115:737-741.
2. King TF, Conway GS. Swyer syndrome. *Curr Opin Endocrinol Diabetes Obes*. 2014;21:504-510.
3. de Vries ALC, Roehle R, Marshall L, Frisén L, van de Grift TC, Kreukels BPC, Bouvattier C, Köhler B, Thyen U, Nordenström A, Rapp M, Cohen-Kettenis PT; dsd-LIFE Group. Mental health of a large group of adults with disorders of sex development in six European countries. *Psychosom Med*. 2019;81:629-640.
4. Hughes IA, Houk C, Ahmed SF, Lee PA. Consensus statement on management of intersex disorders. *Arch Dis Child*. 2006;91:554-563. Epub 2006 Apr 19
5. Şentürk Pılan B, Özbaran B, Çelik D, Özcan T, Özen S, Gökşen D, Ulman İ, Avanoğlu A, Tiryaki S, Onay H, Çoğulu Ö, Özkinay F, Darcan Ş. Quality of life and psychological well-being in children and adolescents with disorders of sex development. *J Clin Res Pediatr Endocrinol*. 2021;13:23-33.
6. McCluskey SE, Lacey JH. Anorexia nervosa in a patient with XY gonadal dysgenesis. *Br J Psychiatry*. 1992;160:114-116.